

# The Impact of Vitamin D Deficiency on Some Immunomodulatory Markers in Pregnant Women with Threatened Abortion

Rafal J. Al-Saigh<sup>1</sup>, Enass Najem Oubaid<sup>2</sup>, Nada Khazal Kadhim Hindi<sup>3</sup>, Ahmed H Al-Humadi<sup>4</sup>, Hussam Wahab Al-Humadi<sup>5\*</sup>

Department of Clinical Laboratory sciences, College of Pharmacy, University of Babylon, Babylon Province, Iraq<sup>1</sup>

Department of Pharmacognosy, College of Pharmacy, University of Babylon, Babylon Province, Iraq<sup>2</sup>  
Department of Basic and Medical Science, College of Nursing, University of Babylon, Babylon Province, Iraq<sup>3</sup>  
Medical School, National and Kapodestrian University of Athens, 75 Mikras Asias str., Athens, Greece<sup>4</sup>  
Dean of College of Pharmacy, Department of Clinical Pharmacy, Babylon University; Babylon Province, Iraq<sup>5</sup>

Corresponding Author: 5\*



## Keywords:

Vitamin D, Pregnancy, Threatened Abortion, IL-6, TNF- $\alpha$ , INF- $\gamma$ .

## ABSTRACT

The protective mechanism of vitamin D (VD) on the immune system played an important role behind many health problems including abortion. Our aim was to investigate the levels of VD and some immunomodulatory markers (IL-6, TNF- $\alpha$  and INF- $\gamma$ ) in pregnant women with threatened abortion (TA) and their effects on their obstetric health. This study including 132 pregnant participants with TA visiting maternity hospital in Babel province, Iraq. The demographic data including previous and current obstetrical history and body mass index were collected by well-trained researcher following structured questionnaire after obtaining their verbal consent. Laboratory Investigations were done with estimation of VD, IL-6, TNF- $\alpha$  and INF- $\gamma$ . The mean level of VD was  $19.83 \pm 43.59$  ng/ml while the mean levels of IL-6:  $389.45 \pm 70.88$ , TNF- $\alpha$ :  $118.78 \pm 44.11$  and INF- $\gamma$ :  $100.39 \pm 20.13$  pg/ml with related percentage among participants for low VD was 84%, high IL-6 was 66%, high TNF- $\alpha$  was in 43% but only 23% had high INF- $\gamma$  in relation with normal values. The frequency of abortion was significantly increased with low VD and high IL-6 ( $p < 0.05$ ) with high correlation between them ( $r^2 = 0.89$ ). High incidence of TA associated with low level of VD accompanied with high serum levels of IL-6. This adverse pregnancy outcome may be associated with some form of autoimmune diseases with evidence that hypovitaminosis D was correlated with high proinflammatory markers (IL-6, TNF- $\alpha$  and INF- $\gamma$ ) and worse form of TA and the role of VD is to correct these pro-inflammatory markers.



This work is licensed under a Creative Commons Attribution Non-Commercial 4.0 International License.

## 1. Introduction

Threatened abortion is a clinical pregnancy associated with an intrauterine viable fetus that commonly accompanied by painless vaginal bleeding and may normally continuation of the pregnancy or be accompanied by pain to become an inevitable abortion [1]. Identifying the associated factors can play an important role in preventing it and improving pregnancy outcomes. There is a crucial evidence for vitamin D in the body's immune system. Most of these researches have been done in cultured cells and in animals to prove the active control of vitamin D metabolism. Vitamin D receptor is found in significant concentrations in the T lymphocyte mostly the mature CD-8 T lymphocytes. This suggests that the action of T lymphocytes proved through calcitriol [2], [3]. The role of vitamin D is to block the effects of adaptive immune system leading to prevent the autoimmunity that proved by animal studies which showed with absent of calcitriol, the immune cells are attacked the healthy cells of the body leading to autoimmunity [2]. Moreover, vitamin D can be blocking the production of pro-inflammatory cytokines like interleukin-6 (IL-6), Tumor Necrotizing Factor- $\alpha$  (TNF- $\alpha$ ) and Interferon- $\gamma$  (IFN- $\gamma$ ) and that may also have associated with stimulating the production of anti-inflammatory cytokines and decreasing in killer T cells production [2- 4]. On other hand, the deficiency of vitamin D can inhibit the production of B lymphocytes leading to blocks the production of immunoglobulins [3- 5].

In normal immune response, T helper-1 cells produce INF- $\gamma$ , IL-2, and TNF- $\alpha$ , and this response is strongly addressed to intracellular pathogens as viruses while T helper-2 lymphocytes produce mainly Tissue Growth Factor (TGF)- $\beta$ 1 and IL-4 and IL-5, and these mediators are linked with extracellular pathogens as bacteria and parasites [6] that proved the same role of VD decreases the production of T helper-1 producers like INF- $\gamma$ , IL-2, and TNF- $\alpha$  and increases the production of T helper-2 producers like IL-4 and IL-5 [6], [7].

Furthermore, the immunomodulating effects of VD and its analogs have been well-characterized in dendritic cells (DCs), which are as antigen presenting cells (APCs). In response to inflammatory signals, VD strongly impairs the migration and maturation of DCs, which reduces antigen presentation capacity and activation of T lymphocytes with marked reduction in pro-inflammatory ILs that contribute as an induction of tolerogenic state [3], [4].

The aim of study was to investigate the levels of VD and some immunomodulatory markers (IL-6, TNF- $\alpha$  and INF- $\gamma$ ) in pregnant women with TA and their effects on their obstetric health.

## 2. SUBJECTS AND METHODS

### 2.1 Patients

The number of patients was 132, study was carried out at maternity hospital in Babel province, Iraq during the period from June, 2019 until March, 2020. Sample size was taken by consecutive manner.

### 2.2 Exclusion criteria

Eligible patients should not document with known chronic liver, kidney, thyroid or gastrointestinal diseases, hypertension, diabetes, or using any medications that affect vitamin D level (e.g. anticonvulsants, antituberculosis drugs, etc)

### 2.3 Criteria of threatened abortion

Persistent vaginal bleeding at first 20 weeks of gestation with viable fetus [1].

#### **2.4 Ethics statement**

The study was carried out in compliance with the Declaration of Helsinki principles and was reviewed and approved by local research ethics committees of Faculty of Pharmacy, University of Babylon (Protocol No. 193/2019)). All-important study material was communicated to the patient group, and they were told that their rejection or withdrawal from the trial would have no impact on the medical care they received. After gaining their verbal consent, the data was collected by a well-trained researcher using a standardized questionnaire.

#### **2.5 Demographic and clinical variables**

Maternal age, residency, socioeconomic states, educational and occupation states, previous and current obstetrical history and body mass index (BMI) were collected by questionnaire to all participants.

The effects of maternal age were investigated using continuous and categorical approaches (coded as age < 25, 25-30, 30-35, and 35-40 and > 40 years).

Medical records were abstracted to a certain their anthropomorphic characteristics as well as their medical status throughout gestation. Pregnancy BMI was calculated using the admission weight and height measurements.

#### **2.6 Laboratory Investigation**

During admission period, 5ml venous blood sample was collected to measure vitamin D, IL-6, TNF- $\alpha$ , and IFN- $\gamma$ .

#### **2.7 Level of Serum vitamin D3 (25-hydroxyvitamin-D)**

Estimation of vitamin D level from blood samples were determined by using a commercial enzyme-linked immunosorbent assay (ELISA) (Demediteq: DE1971, Germany). According to the manufacturer's protocol, each assay was run with known standards (provided with the kit) that were used to determine the quantity of vitamin D in each sample in ng/ml. The range of detection was 3–300 ng/mL. Patients' vitamin D status was evaluated according to vitamin D concentrations into deficient Levels below 25 ng/ml, insufficient between 25–75 ng/ml and normal level  $\geq 75$  ng/ml [8].

#### **2.8 Immunomodulatory markers: IL-6, TNF- $\alpha$ , and IFN- $\gamma$ levels**

Inflammatory cytokines indices were measured by ELISA technique (elabscience company, USA). IL-6 is by E-EL-H0192 kit with normal values ranging from (104.35-238.90 pg/ml), TNF- $\alpha$  by E-EL-H 0109 kit with normal values ranging from (24.10-86.50 pg/ml), and IFN- $\gamma$  by E-EL- H0108 kit with normal values ranging from (51.90-100.10 pg/ml). According to the manufacturer's protocol; each assay was run with known standards (provided with the kit) that were used to determine the quantity of cytokines in each sample.

#### **2.9 Statistical analysis**

To explore relationships and differences in Demographic, clinical factors, VD, and immunomodulatory markers, descriptive measures, one way and two-way ANOVA analysis were utilized. For all analyses, the significance level was fixed at a probability (P) of less than or equal to 0.05. GraphPad Prism 5.3 on Windows was used to conduct all analyses (GraphPad Software, San Diego, CA, USA).

### **3. RESULTS**

Serum VD levels ranged from 2.8 to 68.5 ng/ml with mean level was  $19.83 \pm 43.59$  ng/ml. The distribution

of VD among participant population was 84% (mean:  $16.2 \pm 10.6$  ng/ml) with deficient vitamin D ( $< 25$  ng/ml), 11% (mean:  $32.8 \pm 17.6$ ) with insufficient vitamin D (25-75 ng/ml) and only 5% (mean:  $76.5 \pm 3.6$  ng/ml) with normal vitamin D level ( $> 75$  ng/ml) (Table 1). The rang of IL-6 levels was 133.94-438.90 pg/ml with mean level of  $389.45 \pm 70.88$  and associated with 66% of participant higher than normal rang. The rang of TNF- $\alpha$  levels was 78.51-162.31 pg/ml with mean level of  $118.78 \pm 44.11$  and associated with 43% of participant higher than normal rang. The rang of INF- $\gamma$  levels was 80.22-119.93 pg/ml with mean level of  $100.39 \pm 20.13$  and associated with only 23% of participant higher than normal rang (Table 1).

**Table 1.** Serum Levels of Vitamin D, Interleukin-6 (IL-6), Tumor Necrotizing Factor- $\alpha$  (TNF-  $\alpha$ ) and Interferon- $\gamma$  (IFN-  $\gamma$ ) in Threatened Abortion Participants (n=132).

Immunomodulatory Markers	Means	Std. Deviation	Std. Error	Lower 95% CI of mean	Upper 95% CI of mean
<b>Vitamin D</b>	19.83 ng/ml	43.59	3.794	12.32	27.34
<b>IL-6</b>	389.5 pg/ml	70.88	6.169	377.2	401.7
<b>TNF-<math>\alpha</math></b>	118.8 pg/ml	44.11	3.839	111.2	126.4
<b>INF-<math>\gamma</math></b>	100.4 pg/ml	20.13	1.752	96.92	103.9

The women in the study ranged in age from 18 to 41 years old, with 70% of them being between the ages of 25 and 35. The majority of the participants were from a middle socioeconomic status (40%) with a secondary education (54%) and were housewives (77%) or multipara (1-3 para) (47%).

**Table 2.** sociodemographic data, previous obstetrical and medical information in the studied population in relation to the serum level of vitamin D.

Parameters	Sub-parameters	Normal vitamin D N = 12	Insufficient vitamin D N = 22	Deficient vitamin D N = 98	<i>P value</i>
<b>Age</b>	< 25 yrs	3 (2%)	4 (3%)	20 (15%)	<i>P</i> <0.001
	25-30 yrs	3 (2%)	8 (6%)	60 (45%)	
	31-35 yrs	1 (1%)	3 (2%)	17 (13%)	
	36-40 yrs	0 (0%)	4 (3%)	8 (6%)	
	> 40 yrs	0 (0%)	0 (0%)	1 (1%)	
<b>Socioeconomic state</b>	Low	0 (0%)	4 (3%)	37 (34%)	<i>P</i> <0.001
	Medium	2 (2%)	5 (4%)	42 (39%)	
	High	3 (3%)	4 (4%)	14 (11%)	
<b>Educational Level</b>	Elementary	0 (0%)	4 (3%)	29 (22%)	<i>P</i> <0.001
	Secondary	3 (2%)	8 (6%)	63 (48%)	
	Bachelor or >	4 (3%)	3 (2%)	18 (14%)	
<b>Occupation</b>	Housewife	4 (3%)	11 (8%)	89 (68%)	<i>P</i> <0.001
	Employee	3 (2%)	4 (3%)	21 (16%)	
<b>BMI (kg/m<sup>2</sup>)</b>	<25	0 (0%)	3 (2%)	13 (10%)	<i>P</i> <0.001

	25-30	3 (2%)	3 (2%)	68 (52%)	
	> 30	4 (3%)	10 (8%)	28 (22%)	
<b>Number of Parity</b>	Primiparous	6 (4%)	6 (4%)	42 (32%)	<i>P&lt;0.001</i>
	1-3 Para	1 (1%)	8 (6%)	55 (42%)	
	> 3 Para	0 (0%)	1 (1%)	13 (10%)	
<b>Status of Pregnancy</b>	Singleton	4 (3%)	10 (7%)	87 (66%)	<i>P&lt;0.001</i>
	Multiple	3 (2%)	5 (4%)	23 (17%)	
<b>History of abortion</b>	Yes	1 (1%)	6 (5%)	34 (26%)	<i>P&lt;0.001</i>
	No	5 (4%)	8 (6%)	77 (58%)	
<b>History of Fetal Death</b>	Yes	0 (0%)	3 (2%)	14 (10%)	<i>P&lt;0.001</i>
	No	6 (5%)	12 (9%)	98 (74%)	
<b>Dietary History</b>	Bad	1 (1%)	6 (5%)	73 (55%)	<i>P&lt;0.001</i>
	Good	5 (4%)	8 (6%)	39 (29%)	
<b>Sun exposure</b>	> One hr	3 (2%)	3 (2%)	16 (12%)	<i>P&lt;0.001</i>
	< One hr	4 (3%)	12 (9%)	94 (72%)	


BMI: body mass index

Participants had a 41% chance of having a previous miscarriage, and a 12% chance of having a fetal death. BMI was 56% of the women were between 25-30 kg/m<sup>2</sup>, while 31 % for participant more than 30 kg/m<sup>2</sup> and only 12 % was below 25 kg/m<sup>2</sup>. Regarding the anemia, most of participant was non-anemic (58%) in spite of bad dietary history during pregnancy (59%) (Table 2).

Moreover, there was a significant relation ( $P > 0.001$ ) between the numbers of previous mischarge (Table 3) and the levels of VD (Figure 1) that increase in the significances between the numbers of previous miscarriage in the deficient status of vitamin D ( $P < 0.001$ ).


**Table 3.** The numbers of previous abortions among participant women (n=132).

<b>Previous abortion</b>	<b>0</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4 or more</b>
<b>Numbers</b>	42	34	26	19	11



**Figure 1.** Relation between the levels of vitamin D and the numbers of previous abortion in study participant (n=132). The number of asterisks (\*) corresponds to the level of the statistical significance (\*\*P<0.01, \*\*\*P<0.001).


Furthermore, there was also a significant relation between the numbers of previous miscarriage and the levels of IL-6, TNF- $\alpha$  and INF- $\gamma$  (Figure 2), these significances varied according the numbers of previous miscarriage (significances ranging from P<0.05 to P<0.001). On other hand, the frequency of abortion was significantly increased with low VD and high IL-6 (p<0.05) with high correlation between them (R<sup>2</sup>=0.89) (Figure 3).



**Figure 2.** Relation between the levels of Interleukin-6 (IL-6), Tumor Necrotizing Factor- $\alpha$  (TNF-  $\alpha$ ) and Interferon- $\gamma$  (INF-  $\gamma$ ) and the numbers of previous abortion in study participant (n=132). The number of asterisks (\*) corresponds to the level of the statistical significance (\*P<0.05, \*\*P<0.01, \*\*\*P<0.001).

#### 4. DISCUSSION

There are several physiological alterations associated with pregnancy including changes in VD metabolism making only 16% of women had normal levels of VD [9]. Furthermore, the immune system is also involved in various pathological and physiological events during pregnancy [10].



**Figure 3.** Correlation between the levels of vitamin D levels and Interleukin-6 (IL-6) in relation to

participant (n=132)

The presence of different proteins, cytokines, and cells that can contribute to the development and maintenance of pregnancy can lead to the rejection or even termination of the fetus due to the disruption of its immunomodulation (11). T helper cells are known to be very pathogenic and can trigger autoimmune and inflammatory diseases [12], [13].

Moreover, T helper cells can also contribute to the development and maintenance of pregnancy by regulating the expression of chemokines and proinflammatory cytokines [14]. These cells can also infiltrate and destroy tissue. The presence of cytokines in pregnancy can also alter the effects of different hormones [15].

Some studies revealed that the presence of high levels of proinflammatory cytokines in women who were threatened with an abortion [16], [17] was associated with an increased likelihood of having a negative outcome. These findings support the idea that low levels of these cytokines during early pregnancy are associated with a better chance of having a successful pregnancy [18], [19].

In vitro investigations also shown that participants with spontaneous abortion had higher IL-6 levels than those with normal pregnancy [20], [21], but that IL-6 levels were relatively high in in vitro fertilization produced pregnancies [22]. The release of cytokines into the surrounding fluid by activated T cells, and the rise of these cytokines (IL-6, TNF-, and INF-) in peripheral levels after they were released, suggests immune system activation and may reflect immunologic rejection of the fetus as a cause of spontaneous abortion [23].

Previous research has linked high amounts of IL-6 in pregnant women's peripheral blood and placenta to low levels of VD [24]. VD has the ability to lessen the occurrence of spontaneous combustion [25].

In pregnancy IL-6 plays a crucial role in chorioamnionitis with involving of rupturing the fetal membranes at term [26]. However, TNF- $\alpha$  is believed to play a role in pre-implantation embryo transport by inducing local contractions [27]. No significant evidence that the changing in the level of INF- $\gamma$  may affected the outcome of pregnancy in spite of antiviral and cytotoxic effects of it [28] but according to several research, VD status and considerable secretion of greater amounts of IFN- women with recurrent abortion have a high link with women who have at least one healthy life birth without spontaneous abortions or infertility [29].

Our findings point to a more immediate and subtle role for the immune system in threatening abortion than an eventual inflammatory pathogenesis, which has been identified as an immunomodulator in different mechanisms during pregnancy, including VD.

## 5. CONCLUSIONS

In summary, High incidence of Threatened Abortion associated with low level of Vitamin D accompanied with high serum levels of IL-6. This adverse pregnancy outcome may be associated with some form of autoimmune diseases. Hypovitaminosis D was correlated with high proinflammatory markers (IL-6, TNF-  $\alpha$  and IFN-  $\gamma$ ) and worse form of Threatened Abortion and the role of Vitamin D is to correct these pro-inflammatory markers.

## 6. CONFLICT OF INTEREST



No conflict of interest

## 7. ACKNOWLEDGEMENT

Thanks to Babylon Health Office, Maternity and Child hospital in Babel province, Iraq for their precious help and assistance.

## 8. AUTHOR CONTRIBUTION

All authors shared in conceptualization with direct and intellectual contribution to the work, writing, reference search, table and figures preparation and approved it for publication.

## 9. REFERENCES

- [1] ACOG Committee on Practice Bulletins--Gynecology. ACOG Practice Bulletin: No 15: Premenstrual syndrome. *Obstet Gynecol.* 2000 Apr;95(4):suppl 1-9. Retraction in: *Obstet Gynecol.* 2012 Aug;120(2 Pt 1):405. PMID: 24829972.
- [2] von Essen MR, Kongsbak M, Schjerling P, Olgaard K, Odum N, Geisler C. Vitamin D controls T cell antigen receptor signaling and activation of human T cells. *Nat Immunol.* 2010 Apr;11(4):344–349.
- [3] Deluca HF, Cantorna MT. Vitamin D: its role and uses in immunology. *FASEB J.* 2001 Dec;15(14):2579–2585.
- [4] Hewison M. An update on vitamin D and human immunity. *Clin Endocrinol (Oxf).* 2012 Mar;76(3):315–325.
- [5] Aranow C. Vitamin D and the immune system. *J Investig Med.* 2011 Aug;59(6):881–886.
- [6] Cantorna MT, Zhu Y, Froicu M, Wittke A. Vitamin D status, 1,25-dihydroxyvitamin D3, and the immune system. *Am J Clin Nutr.* 2004 Dec;80(6 Suppl):1717S–1720S.
- [7] Mahon BD, Wittke A, Weaver V, Cantorna MT. The targets of vitamin D depend on the differentiation and activation status of CD4 positive T cells. *J Cell Biochem.* 2003 Aug 1;89(5):922–932.
- [8] López-Vicente, M., Sunyer, J., Lertxundi, N. et al. Maternal circulating Vitamin D3 levels during pregnancy and behaviour across childhood. *Sci Rep* 9, 14792 (2019). <https://doi.org/10.1038/s41598-019-51325-3>
- [9] Wei SQ, Qi HP, Luo ZC, et al. Maternal Vitamin D status and adverse pregnancy outcomes: a systematic review and meta-analysis. *J Matern Fetal Neonatal Med* 2013;26:889-99.
- [10] Regal JF, Gilbert JS, Burwick RM. The complement system and adverse pregnancy outcomes. *Mol Immunol.* 2015 Sep;67(1):56-70. doi: 10.1016/j.molimm.2015.02.030. Epub 2015 Mar 21. PMID: 25802092; PMCID: PMC4447554.
- [11] Kwak-Kim J, Bao S, Lee SK, Kim JW, Gilman-Sachs A. Immunological modes of pregnancy loss: inflammation, immune effectors, and stress. *Am J Reprod Immunol.* 2014 Aug;72(2):129- 40. doi: 10.1111/aji.12234. Epub 2014 Mar 24. PMID: 24661472.

- [12] Zhang XM, Liu CY, Shao ZH. Advances in the role of helper T cells in autoimmune diseases. *Chin Med J (Engl)*. 2020 Apr 20;133(8):968-974. doi: 10.1097/CM9.0000000000000748. PMID: 32187054; PMCID: PMC7176439.
- [13] Kunkl M, Frasca S, Amormino C, Volpe E, Tuosto L. T Helper Cells: The Modulators of Inflammation in Multiple Sclerosis. *Cells*. 2020 Feb 19;9(2):482. doi: 10.3390/cells9020482. PMID: 32093011; PMCID: PMC7072830.
- [14] Zhang JM, An J. Cytokines, inflammation, and pain. *Int Anesthesiol Clin*. 2007 Spring;45(2):27-37. doi: 10.1097/AIA.0b013e318034194e. PMID: 17426506; PMCID: PMC2785020.
- [15] Guzeloglu-Kayisli O, Kayisli UA, Taylor HS. The role of growth factors and cytokines during implantation: endocrine and paracrine interactions. *Semin Reprod Med*. 2009 Jan;27(1):62-79. doi: 10.1055/s-0028-1108011. Epub 2009 Feb 5. PMID: 19197806; PMCID: PMC3107839.
- [16] Freis A, Schlegel J, Daniel V, Jauckus J, Strowitzki T, Germeyer A. Cytokines in relation to hCG are significantly altered in asymptomatic women with miscarriage - a pilot study. *Reprod Biol Endocrinol*. 2018 Sep 28;16(1):93. doi: 10.1186/s12958-018-0411-5. PMID: 30266090; PMCID: PMC6162891.
- [17] Paradisi R, Porcu E, Venturoli S, Maldini-Casadei M, Boni P. Maternal serum levels of pro-inflammatory cytokines in missed and threatened abortion. *Am J Reprod Immunol*. 2003 Oct;50(4):302-8. doi: 10.1034/j.1600-0897.2003.00065.x. PMID: 14672332.
- [18] Yockey LJ, Iwasaki A. Interferons and Proinflammatory Cytokines in Pregnancy and Fetal Development. *Immunity*. 2018 Sep 18;49(3):397-412. doi: 10.1016/j.immuni.2018.07.017. PMID: 30231982; PMCID: PMC6152841.
- [19] Chatterjee P, Chiasson VL, Bounds KR, Mitchell BM. Regulation of the Anti-Inflammatory Cytokines Interleukin-4 and Interleukin-10 during Pregnancy. *Front Immunol*. 2014 May 27;5:253. doi: 10.3389/fimmu.2014.00253. PMID: 24904596; PMCID: PMC4034149.
- [20] Liu RX, Wang Y, Wen LH. Relationship between cytokine gene polymorphisms and recurrent spontaneous abortion. *Int J Clin Exp Med*. 2015 Jun 15;8(6):9786-92. PMID: 26309657; PMCID: PMC4538128.
- [21] Koumantaki Y, Matalliotakis I, Sifakis S, Kyriakou D, Neonaki M, Goymenou A, Koumantakis E. Detection of interleukin-6, interleukin-8, and interleukin-11 in plasma from women with spontaneous abortion. *Eur J Obstet Gynecol Reprod Biol*. 2001 Sep;98(1):66-71. doi: 10.1016/s0301-2115(01)00289-5. PMID: 11516802.
- [22] Yang J, Yang X, Yang H, Bai Y, Zha H, Jiang F, Meng Y. Interleukin 6 in follicular fluid reduces embryo fragmentation and improves the clinical pregnancy rate. *J Assist Reprod Genet*. 2020 May;37(5):1171-1176. doi: 10.1007/s10815-020-01737-2. Epub 2020 Mar 18. PMID: 32189182; PMCID: PMC7244683.
- [23] Chatterjee P, Chiasson VL, Bounds KR, Mitchell BM. Regulation of the Anti-Inflammatory Cytokines Interleukin-4 and Interleukin-10 during Pregnancy. *Front Immunol*. 2014 May 27;5:253. doi:

10.3389/fimmu.2014.00253. PMID: 24904596; PMCID: PMC4034149.

[24] Kim H, Baek S, Hong SM, Lee J, Jung SM, Lee J, Cho M, Kwok SK, Park SH. 1,25-dihydroxy Vitamin D3 and Interleukin-6 Blockade Synergistically Regulate Rheumatoid Arthritis by Suppressing Interleukin-17 Production and Osteoclastogenesis. *J Korean Med Sci.* 2020 Feb 17;35(6):e40. doi: 10.3346/jkms.2020.35.e40. PMID: 32056400; PMCID: PMC7025907.

[25] Schröder-Heurich B, Springer CJP, von Versen-Höynck F. Vitamin D Effects on the Immune System from Periconception through Pregnancy. *Nutrients.* 2020 May 15;12(5):1432. doi: 10.3390/nu12051432. PMID: 32429162; PMCID: PMC7284509.

[26] Omere C, Richardson L, Saade GR, Bonney EA, Kechichian T, Menon R. Interleukin (IL)-6: A Friend or Foe of Pregnancy and Parturition? Evidence From Functional Studies in Fetal Membrane Cells. *Front Physiol.* 2020 Jul 24;11:891. doi: 10.3389/fphys.2020.00891. PMID: 32848846; PMCID: PMC7397758.

[27] Neira JA, Tainturier D, Peña MA, Martal J. Effect of the association of IGF-I, IGF-II, bFGF, TGF-beta1, GM-CSF, and LIF on the development of bovine embryos produced in vitro. *Theriogenology.* 2010 Mar 15;73(5):595-604. doi: 10.1016/j.theriogenology.2009.10.015. Epub 2009 Dec 24. PMID: 20035987.

[28] Yockey LJ, Iwasaki A. Interferons and Proinflammatory Cytokines in Pregnancy and Fetal Development. *Immunity.* 2018 Sep 18;49(3):397-412. doi: 10.1016/j.immuni.2018.07.017. PMID: 30231982; PMCID: PMC6152841.

[29] Tavakoli M, Jeddi-Tehrani M, Salek-Moghaddam A, Rajaei S, Mohammadzadeh A, Sheikhhasani S, Kazemi-Sefat GE, Zarnani AH. Effects of 1,25(OH)<sub>2</sub> vitamin D<sub>3</sub> on cytokine production by endometrial cells of women with recurrent spontaneous abortion. *Fertil Steril.* 2011 Sep;96(3):751-7. doi: 10.1016/j.fertnstert.2011.06.075. PMID: 21880282.